

Benefits/Burdens of Tube Feeding/PEG Placement for Adults

	Dysphagic Stroke (Patients with previous good quality of life, high functional status ¹ and minimal co-morbidities)	Dysphagic Stroke (Patients with decreased level of consciousness, multiple co-morbidities, poor functional status ¹ prior to CVA)	Neurodegenerative Disease [e.g., Amyotrophic Lateral Sclerosis (ALS)]	Persistent Vegetative State (PVS)	Frailty (Patients with multiple co-morbidities, poor functional status, failure to thrive and pressure ulcers ²)	Advanced Dementia (Patients needing help with daily care, having trouble communicating, and/or incontinent)	Advanced Cancer (Excludes patients with early stage esophageal & oral cancer)	Advanced Organ Failure (Patients with CHF, renal or liver failure, COPD, anorexia-cachexia syndrome)
Prolongs Life	<i>Likely</i>	<i>Likely in the short term</i> <i>Not likely in the long term</i>	<i>Likely</i>	<i>Likely</i>	Not Likely	Not Likely	Not Likely	Not Likely
Improves Quality of Life and/or Functional Status	up to 25% regain swallowing capabilities	Not Likely	Uncertain	Not Likely	Not Likely	Not Likely	Not Likely	Not Likely
Enables Potentially Curative Therapy/Reverses the Disease Process	Not Likely	Not Likely	Not Likely	Not Likely	Not Likely	Not Likely	Not Likely	Not Likely

This grid reflects only certain conditions. Some examples of other conditions where direct enteral feeding would be indicated include radical neck dissections, esophageal stenosis and motility diseases, post intra-thoracic esophageal surgery and safer nutrition when the alternative would be parenteral hyperalimentation.

Benefits of PEG placement rather than feeding orally:

- For dysphagic stroke patients in previous good health, patients with ALS, and patients in a persistent vegetative state, may prolong life
- For dysphagic stroke patients in previous poor health, may prolong life in the short-term (days to weeks)
- Enables family members/caregivers to maintain hope for future improvement
- Enables family members/caregivers to avoid guilt/conflict associate with choosing other treatment options
- Allows family/caregivers additional time to adjust to possibility of impending death

Burdens of PEG placement rather than feeding orally:

- 75% of stroke patients previously in good health not likely to have improved quality of life and/or functional status
- PVS patients not likely to have improved quality of life and/or functional status
- Possible patient agitation resulting in use of restraints
- Risk of aspiration pneumonia is the same or greater than that of patient being handfed
- Stroke patients previously in poor health, frail patients, and patients w/advanced dementia, cancer or organ failure have been reported to experience side effects: PEG site irritation or leaking (21%), diarrhea (22%), nausea (13%) and vomiting (20%)

This information is based predominately on a consensus of current expert opinion. It is not exhaustive. There are always patients who prove exceptions to the rule.

1. Functional Status refers to Activities of Daily Living. For more information on the CFS visit http://geriatricresearch.medicine.dal.ca/clinical_frailty_scale.htm A poor functional status means full or partial dependency in bathing, dressing, toileting, feeding, ambulation, or transfers.
2. Matched residents with and without a PEG insertion showed comparable sociodemographic characteristic, rates of feeding tube risk factors, and mortality. Adjusted for risk factors, hospitalized NH residents receiving a PEG tube were 2.27 times more likely to develop a new pressure ulcer (95% CI, 1.95-2.65). Conversely, those with a pressure ulcer were less likely to have the ulcer heal when they had a PEG tube inserted (OR 0.70 [95% CI, 0.55-0.89]). Teno JM, Gozalo P, Mitchell SL, Kuo S, Fulton AT, Mor V. Feeding Tubes and the Prevention or Healing of Pressure Ulcers. *Archives of internal medicine*. 2012;172(9):697-701. doi:10.1001/archinternmed.2012.1200.
3. Callahan CM, Haag KM, Weinberger M, et.al. Outcomes of Percutaneous Endoscopic Gastrostomy among Older Adults in a Community Setting. *J Am Geriatr Soc*. 2000 Sep; 48(9):1048-5

Benefits of feeding orally rather than inserting a PEG:

- Patient able to enjoy the taste of food
- Patient has greater opportunity for social interaction
- Patient's wishes and circumstances can be taken into consideration as pertains to pace, timing and volume of feeding

Burdens of feeding orally rather than inserting a PEG:

- Requires longer period of time to feed a patient
- Patient/family worry about "not doing everything in their power" to address the feeding problem and/or "starving patient"
- Patient/family feel that in not choosing option that could possibly prolong life, they are hastening death


Benefits/Burdens of Tube Feeding/PEG Placement for Adults

Legal and Ethical Issues

<p>For patients who CAN make decisions for themselves,</p>	<p>Usual standards of informed consent (or refusal) apply. This applies to persons with developmental disabilities who can decide.</p>	<p>Like with any other procedure, the physician will discuss the pros and cons of a feeding tube with the patient, and, if clinically indicated, the patient can agree to have one or not. If they choose to have a feeding tube at one point in time, they can choose to withdraw it at a later date if it is no longer meeting their goals or needs (provided they still have decision-making capacity). If a patient chooses not to have a feeding tube, food and fluids are offered as tolerated using careful hand feeding.</p>						
<p>For patients who CANNOT make decisions for themselves,</p>	<p>Patient has completed a health care proxy form or has the ability to choose a health care agent. This applies to persons with developmental disabilities.</p>	<p>Formally choosing someone to serve as his/her health care “agent”, the agent is required to make decisions for the patient according to what is known about the patient’s wishes, or, if unknown, according to the patient’s best interests. The agent can make all end-of-life decisions on the patient’s behalf, but the decision must be based on “reasonable knowledge” of the patient’s wishes in the case of withdrawing or withholding of tube feeding. For this reason, it is helpful for the signed health care proxy form to include a statement indicating that conversations have occurred between the patient and the health care agent about artificial hydration and nutrition (tube feeding).</p>						
	<p>Patient has not completed a health care proxy form and the person does not have developmental disability.</p>	<p>The legal standard for withholding or withdrawing a feeding tube is currently different depending upon whether the patient resides in a medical facility (hospital or nursing home) or community (e.g. patient’s home, assisted living facility, etc.). As of September 2011 under New York State law “hospital” means a general hospital or hospice.</p>						
		<p>If such a patient is in hospital or nursing home, New York State law allows for surrogate decision makers to make decisions about tube feeding based upon “substituted judgment” (what is known about, the patient’s wishes), or if unknown based on the patient’s best interests. Surrogate decision makers in NYS are in order of priority:</p> <table border="0" style="width: 100%;"> <tr> <td style="width: 50%;">1. A patient’s authorized guardian</td> <td style="width: 50%;">4. Parent</td> </tr> <tr> <td>2. Spouse (if not legally separated) or domestic partner</td> <td>5. Brother/sister (age 18 or over)</td> </tr> <tr> <td>3. Son or daughter over the age of 18 close friend)</td> <td>6. Close friend (Must complete a signed statement as a close friend)</td> </tr> </table> <p>Additionally, under this circumstance, two physicians must concur that either:</p> <ol style="list-style-type: none"> i. the patient has an illness or injury expected to cause death within six months, or ii. the patient is permanently unconscious, or iii. treatment is inhumane or extraordinarily burdensome and the patient has an irreversible or incurable condition <p>Special requirements exist for an Ethics Review Committee to determine that patient-centered and clinical standards are met:</p> <ol style="list-style-type: none"> i. In a hospital, <u>other than a hospice</u>, if the attending physician disagrees with a decision to withhold or withdraw a feeding tube ii. In a nursing home, for all life-sustaining treatment, including a feeding tube, if the clinical standard that the patient meets is “treatment is inhumane or extraordinarily burdensome and the patient has an irreversible or incurable condition”. 	1. A patient’s authorized guardian	4. Parent	2. Spouse (if not legally separated) or domestic partner	5. Brother/sister (age 18 or over)	3. Son or daughter over the age of 18 close friend)	6. Close friend (Must complete a signed statement as a close friend)
1. A patient’s authorized guardian	4. Parent							
2. Spouse (if not legally separated) or domestic partner	5. Brother/sister (age 18 or over)							
3. Son or daughter over the age of 18 close friend)	6. Close friend (Must complete a signed statement as a close friend)							
		<p>If the patient in a hospital or nursing home has not completed a health care proxy form and no surrogate from the list is available, decisions about withholding or withdrawing tube feeding can be made if two physicians concur that:</p> <ol style="list-style-type: none"> i. life sustaining treatment offers no medical benefit and the patient will die imminently even if treatment is provided, AND ii. the provision of life sustaining treatment would violate accepted medical standards 						
		<p>If the patient has not completed a health care proxy form and is not in hospital or nursing home, the legal standard for making a decision about withholding or withdrawing of feeding tubes is “clear and convincing evidence” of the patient’s wishes. A prior written statement about feeding tubes or artificial nutrition in a Living Will, completion of the New York State Medical Orders for Life Sustaining Treatment (MOLST) or clear prior oral statements by the patient about his or her wishes may provide “clear and convincing evidence.”</p>						
	<p>Patient has not completed a health care proxy form, does not have the ability to choose a health care agent and the person has developmental disabilities.</p>	<p>Physicians must follow the § 1750-b process as outlined on the MOLST Legal Requirements Checklist for Individuals with Developmental Disabilities. Two physicians must determine to a reasonable degree of medical certainty that both of the following conditions are met: (1) the individual has one of the following medical conditions: a. a terminal condition; (briefly describe); or b. permanent unconsciousness; or c. a medical condition other than DD which requires LST, is irreversible and which will continue indefinitely (briefly describe) AND (2) the LST would impose an extraordinary burden on the individual in light of: a. the person’s medical condition other than DD (briefly explain) and b. the expected outcome of the LST, notwithstanding the person’s DD (briefly explain.) If the 1750-b surrogate has requested that artificially provided nutrition or hydration be withdrawn or withheld, one of the following additional factors must also be met: a. there is no reasonable hope of maintaining life (explain); or b. the artificially provided nutrition or hydration poses an extraordinary burden (explain.)</p>						

Approach to Adult Patient Unable to Maintain Nutrition

*Numbers refer to Flow Chart Reference Sheet on page 4



Guidelines are intended to be flexible. They serve as reference points or recommendations, not rigid criteria. Guidelines should be followed in most cases, but there is an understanding that, depending on the patient, the setting, the circumstances, or other factors, care can and should be tailored to fit individual needs.

Flow Chart Reference Sheet

1. Triggers for Assessment of Eating/Feeding/Nutrition:

- Weight loss
- Decreased eating (> 25 % left uneaten after most meals) NOTE: Clinicians often overestimate % eaten
- Pressure ulcers
- Presence of enteral or parenteral feedings
- Apparent aspiration and/or dysphagia following, or in the setting of acute illness

2. Global Assessment:

➤ Assess Parameters of Nutritional Status

- Weight change (1-2% or more in 1 week, 5% or more in one month, 7.5% or more in 3 months, 10% or > in 6 months)
- Account for possible fluid imbalance
- Body mass index of < 18.5 KG/meter squared
- Abnormal lab tests (albumin, pre-albumin, cholesterol, lymphocyte count)

➤ Identify Factors that Impede Ability to Take In Adequate Amounts of Food

- Physical limitations, visual problems
- Chewing problems (problems with mouth, teeth, dentures)
- Swallowing problems (feeding position, consistencies, bolus size, conducive environment, stimulus to swallow: verbal and tactile)
- Conditions that decrease nutrient intake (nausea, vomiting, constipation, cancer, shortness of breath, weakness)
- Alterations in taste secondary to medications, dry mouth, food options

➤ Identify Additional Problems in Relation to Nutritional Status

- Mental (dementia, depression, anxiety, delusions, apathy)
- Communication problems resulting in inability to make needs known
- Neurologic conditions

➤ Perform Medical Assessment

- Stage of illness, prognosis, pain
- Assess for constipation/fecal impaction
- Adverse medication effects
 - Address the use of medications that can adversely affect either the ability to eat or the desire to do so. Classes of such drugs include those that induce dry mouth, decrease attentiveness, provoke movement disorders and/or cause GI distress of esophagitis.
 - Specific drugs might include:
 - ❖ sedatives: lorazepam; clonazepam, etc.
 - ❖ antipsychotics: risperidone, quetiapine, aripiprazole, etc.
 - ❖ cholinergic drugs for Alzheimer's: donepezil, galantamine, rivastigmine; anticholinergics: tolterodine, oxybutynin chloride
 - ❖ GI irritants or anorexigenics: NSAIDs, COX II, bisphosphonates, opioids, digoxin, theophylline, antibiotics, iron, calcium, memantine, SSRIs

➤ Assess Hydration Status

- Urine output
- Orthostatic hypotension

3. Assessment of Knowledge, Values and Goals:

➤ Conversation with Relevant Individuals Should Include Discussion of

- their understanding of current illness, health status, functional ability
- advance directives or what the patient/individual would want if able to communicate
- hopes and concerns about future course of illness
- patient values, preferences, cultural and spiritual concerns
- general goals for care (not technical options)
- all viable options for addressing nutritional problems
- placement, operation, care required of PEG
- for particular condition, proven benefits and burdens (and the likelihood of both) of placing PEG (**see Benefits and Burdens grid on page 5**)

4. Discuss/Describe Components of PEG:

➤ Discuss Time-limited Trial with Specific Goals for Care

- Return to baseline level of consciousness
- Acceptable level of functioning
- Nutritional bridge during an acute illness
- Weight gain
- Healing of pressure sores
- Improved biochemical markers of nutrition

➤ With All Appropriate Individuals, Discuss Who (and at What Interval) Will Revisit the Decision to Continue the PEG

5. Careful Attention to Comfort Care is Critical:

- Offer and assist eating if needed but do not force food
- Patient preference should determine type and amount of food
- Excellent mouth care is important.

6. MOLST (Medical Orders for Life-Sustaining Treatment):

- MOLST is a clinical process designed to facilitate communication between health care professionals and patients with advanced illness (or their Health Care Agent or Public Health Law Surrogate or § 1750-b Surrogate) that facilitates shared informed medical decision-making. The result is a set of portable medical orders documented on a bright pink MOLST form that is applicable in all settings and across care transitions, is reviewable, and respects the patient's goals for care regarding the use of cardiopulmonary resuscitation, intubation and mechanical ventilation, hospitalization, feeding tubes and other life-sustaining interventions. To learn more about MOLST, visit CompassionAndSupport.org.

Benefits/Burdens of Tube Feeding/PEG Placement for Adults

Key Recommendations

- Tube feeding decisions should be guided by principles of shared decision-making:
 1. **Begin** by identifying that there is a feeding tube decision to be made
 2. **ASK** – about the patient and family views and experience with current feeding and feeding tubes
 - a. clarify values and preferences
 - b. explore prior experiences
 - c. understand the meaning of the main approaches to the patient and family
 - d. share and explore any prior knowledge of the patient’s past or present wishes in this regard
 3. **TELL** - the clinician then shares his or her experience, including
 - a. medical knowledge about the patient’s disease process and prognosis
 - b. the ability of a feeding tube to alter the disease process and prognosis
 - c. the alternative approach to tube feeding
 - d. any knowledge of the patient’s past values if incapacitated
 4. **ASK** – if there is information that needs more clarification, or if patient/family have questions about what they have heard
 5. **ASK** - the patient and family should be asked what they think the best approach is given information above
 - a. if the clinician agrees, he or she should let it be known that they agree
 - b. if the clinician disagrees, he or she might ask them to say how they came to that conclusion
 6. **ASK** – if the family would like to know what the clinician would recommend based on his or her medical knowledge and knowledge of the patient as a person
 - a. the clinician should ask if they would like to hear what he or she recommends
 - b. if they want to know, the clinician should make a recommendation based on medical knowledge about feeding tubes in the patient’s medical circumstances in light of what is known about the patient’s values
 - c. if they do not want to know, the clinician should offer to talk to them about it in the future if they would like to know
 - d. differences of opinion should be respectfully and carefully negotiated looking for common ground
 - e. a final decision should be postponed until common ground is established (if differences seem unresolvable, then consider a palliative care or ethics consult)
- If tube feeding is elected, develop specific goals and time intervals for reviewing whether these goals have been achieved.
- Physicians should consider the following questions before ordering a swallowing evaluation:
 1. How was the patient swallowing before the hospitalization?
 2. Will the swallow evaluation make a difference for the patient and change the treatment plan?
- Clinicians providing a swallowing evaluation should consider addressing the following questions for follow-up discussions with physicians:
 1. What recommendations do you have about lessening the risk of aspiration while feeding?
 2. What recommendations do you have about food preparation to lessen aspiration risk?
 3. What recommendations do you have about feeding technique to lessen the risk of aspiration?
 4. Will tube feeding in any way lessen the risk of aspiration during natural feeding?
 5. What recommendations do you have about making feeding as enjoyable as possible for the patient?

Benefits/Burdens of Tube Feeding/PEG Placement for Adults

Resources for Physicians

American Geriatrics Society

- [Position Statement](#) - Feeding Tubes in Advanced Dementia (2014)

Choosing Wisely®

Choosing Wisely® is an initiative of the American Board of Internal Medicine Foundation to help clinicians and patients engage in conversations to reduce overuse of tests and procedures, and support patients in their efforts to make smart and effective care choices. More than 70 [specialty society partners](#) have released recommendations with the intention of facilitating wise decisions about the most appropriate care based on a patients' individual situation.

- The [American Academy of Hospice and Palliative Medicine](#) and the [American Geriatric Society](#) listed this recommendation as #1 of their top 5 in 2013, the first year of the Choose Wisely Campaign:

***Don't RECOMMEND percutaneous feeding tubes in individuals with advanced dementia.
Instead, offer oral assisted feedings.***

- The [American Medical Directors Association](#) – The Society for Post-Acute and Long-Term Care Medicine (AMDA) listed this recommendation as #1 of their top 5 in 2013, the first year of the Choose Wisely Campaign:

***Don't INSERT percutaneous feeding tubes in individuals with advanced dementia.
Instead, offer oral assisted feedings.***

Clinical Frailty Scale (CFS)©

Geriatric Medicine Research, Dalhousie University, Halifax, Canada. Permission granted to copy the Clinical Frailty Scale for research and education purposes only.

For more information on the CFS: http://geriatricresearch.medicine.dal.ca/clinical_frailty_scale.htm

CompassionAndSupport.org

Educates and empowers patients, families and professionals on issues related to advance care planning, health care proxies, MOLST (Medical Orders for Life Sustaining Treatment), palliative care, and pain management.

- [Feeding Tubes & Artificial Nutrition & Hydration](#) - addresses the benefits, burdens and challenges of artificial hydration/ nutrition and feeding tubes.
- [MOLST for Professionals](#)

OPWDD MOLST Legal Requirements Checklist for Individuals with Developmental Disabilities

https://opwdd.ny.gov/opwdd_resources/information_for_clinicians/documents/molst-checklist

Professional Society Statements

American Gastroenterological Association. American Gastroenterological Association Medical Position Statement: Guidelines for the Use of Enteral Nutrition. *Gastroenterology*. 1995;108:1280-1301
[http://www.gastrojournal.org/article/0016-5085\(95\)90230-9/pdf](http://www.gastrojournal.org/article/0016-5085(95)90230-9/pdf)

American Geriatrics Society Feeding Tubes in Advanced Dementia Position Statement. American Geriatrics Society Ethics Committee and Clinical Practice and Models of Care Committee Disclosures. *J Am Geriatr Soc*. 2014;62(8):1590-1593.

American Academy of Hospice and Palliative Medicine. Five Things Physicians and Patients Should Question. *Choosing Wisely®* is an initiative of the ABIM Foundation. 2013 Feb. <http://www.choosingwisely.org/societies/american-academy-of-hospice-and-palliative-medicine/>

American Medical Directors Association (AMDA). Altered nutritional status in the long-term care setting. Columbia (MD): American Medical Directors Association (AMDA).2010. 35 p. <http://www.guideline.gov/content.aspx?id=32490>

American Medical Directors Association. Ten Things Physicians and Patients Should Question. *Choosing Wisely®* is an initiative of the ABIM Foundation. 2013 September (1-5) and 2015 March (6-10); rationale for #8 updated 2015 July <http://www.choosingwisely.org/societies/amda-the-society-for-post-acute-and-long-term-care-medicine/>

Benefits/Burdens of Tube Feeding/PEG Placement for Adults

Resources for Patients

ALS Association

Provides information about feeding tubes for people with ALS and caregivers.

Choosing Wisely®

Choosing Wisely® is an initiative of the American Board of Internal Medicine Foundation to help physicians and patients engage in conversations to help make wise decisions about the most appropriate care based on a patients' individual situation. Consumer Reports is a partner in this effort and works with more than 70 specialty societies to create patient educational material about what care is best for them and the right questions to ask their physicians.

- Information about feeding tubes for people with Alzheimer's. Developed in cooperation with the American Academy of Hospice and Palliative Medicine and the American Geriatrics Society.
 - English version
 - Spanish version

CompassionAndSupport.org

Educates and empowers patients, families and professionals on issues related to advance care planning, health care proxies, MOLST, palliative care, and pain management). Videos available:

- Discussing Feeding Tubes & Artificial Nutrition & Hydration - addresses the benefits, burdens & challenges of artificial hydration & nutrition and feeding tubes.
- Patient and Family Stories - describes decision regarding a feeding tube initially inconsistent with the patient's values)

Health in Aging Foundation (Foundation of the American Geriatric Society)

Recommendations on the use of feeding tubes in people with advanced dementia.

University of North Carolina Palliative Care Program

- Making Choices: Feeding Options for Patients with Dementia - Decision aid that explains feeding options for people with dementia. (PowerPoint)
- Helps educate families about feeding problems in people with dementia (video)

Benefits/Burdens of Tube Feeding/PEG Placement for Adults

Recursos para pacientes

[ALS Association](#)

Esta asociación facilita información sobre las sondas de alimentación a las personas con esclerosis lateral amiotrófica (ELA o ALS, por sus siglas en inglés) y a sus cuidadores.

[Choosing Wisely®](#)

Choosing Wisely® es una iniciativa lanzada por el American Board of Internal Medicine Foundation (Panel Estadounidense de la Fundación de Medicina Interna) con el objetivo de propiciar la comunicación entre los médicos y los pacientes a fin de que tomen decisiones acertadas con respecto a la asistencia médica más adecuada para cada paciente en función de la situación particular. Consumer Reports colabora en esta iniciativa y trabaja con más de 70 sociedades especializadas con el objeto de crear materiales educativos para los pacientes sobre la atención médica que más se ajusta a sus necesidades y sobre las preguntas que pueden hacer a los médicos.

- Información sobre las sondas de alimentación para personas con la enfermedad de Alzheimer. Documento elaborado en cooperación con la American Academy of Hospice and Palliative Medicine (Academia Estadounidense de Hospicio y Medicina Paliativa) y la American Geriatrics Society (Sociedad Estadounidense de Geriatría).

Versión en [inglés](#)

Versión en [español](#)

[CompassionAndSupport.org](#)

Este sitio web instruye e informa a los pacientes, familias y profesionales sobre aspectos relacionados con la planificación anticipada de cuidados, los representantes para cuidados médicos, la Orden Médica de Tratamiento para Mantener la Vida (MOLST, por sus siglas en inglés), los cuidados paliativos y el tratamiento del dolor. Videos disponibles:

- [Las sondas de alimentación y la hidratación y nutrición artificiales](#): aborda los beneficios, las cargas y los desafíos de la hidratación y nutrición artificiales y de las sondas de alimentación.
- [Historias de pacientes y familias](#): describe el proceso de toma de decisiones cuando la alimentación por sonda va en contra de los deseos del paciente.

[Health in Aging Foundation](#) (Fundación de la Sociedad Estadounidense de Geriatría)

Esta fundación ofrece recomendaciones sobre el uso de las sondas de alimentación para personas con demencia avanzada.

Programa de cuidados paliativos de la Universidad de Carolina del Norte

- [Toma de decisiones: opciones de alimentación para los pacientes con demencia](#): presentación de ayuda para la toma de decisiones que explica las opciones de alimentación disponibles para las personas con demencia (PowerPoint).
- [Información para las familias sobre los problemas de alimentación de las personas con demencia](#) (video).

Benefits/Burdens of Tube Feeding/PEG Placement for Adults

Patient/Family/Clinician Information To Help You Make a Decision About Tube Feeding/PEG Placement

Problems Swallowing/Eating

People who have a serious illness or are weak may sometimes have problems getting the nourishment we think they need for their body to function properly. Eating and/or swallowing become difficult. When this occurs, the doctor will try to find out what is causing the problem. If treatment or changes in the environment can be made to address this problem, the doctor will see that these changes are made. If the problem cannot be addressed through these changes, the doctor will likely talk to the person and his or her family about tube feeding. One tube feed procedure involves placement of a PEG tube, a feeding tube placed through the skin into the stomach through a small hole in the abdomen.

Discussions with the Doctor about Tube Feeding

Before discussing tube feeding fully, the doctor will ask the person who is sick (or their loved ones if that person cannot make a decision for themselves) about whether or not tube feeding is a procedure that they might be interested in. Some people have very strong feelings about tube feeding and often they have discussed their feelings with loved ones. The doctor will ask about whether the sick person has done any advance care planning whether they have completed a health care proxy or living will. The doctor will ask if the person has had any prior discussions with loved ones about health care preferences in situations like these. Making a decision about tube feeding is often a difficult decision.

If you are making this decision for your loved one, it is important to distinguish what it is they would want for themselves if they could decide for themselves, and to separate that wish from what you would wish for them. It is their wish that should form the basis of the decision.

The Tube Feeding Decision

There are many aspects that need to be considered when making a decision about tube feeding. It is important to consider the advantages, disadvantages and other considerations of feeding tube placement. It is also important to look at the advantages, disadvantages and other considerations to continuation of hand feeding.

The questions you might ask in regards to this decision are:

- Will my loved one live longer, or possibly die sooner, as a result of having a PEG tube placed?
- Will the quality of their life improve and will that quality of life be something they would value?
- Will placing a PEG allow for treatment that is likely to cure their underlying illness?

For example, using tube feeding for a person who had a stroke but was in good health prior to having it will lead to different results than using a tube feeding for a person who has Alzheimer's disease.

Emotions often play a large role in the decision to tube feed. Feelings of guilt about "not doing everything in your power" to help the person and pressure from others may affect the decision making process. Finally, personal beliefs regarding tube feeding influence the decision as well. Health care spokespersons and family members have many questions to consider in making a decision about tube feeding. People who choose not to have tube feedings can be kept comfortable with small sips of liquid and lubrication of their mouths and lips. Most patients will not experience greater comfort because of tube feedings being started. Exceptions to this include some patients with acute injuries that impair their ability to swallow and some people with early cancers of the head and neck and esophagus.

If you and your family members have conflicting views about whether or not the person should have a tube feed placed, it is important to ask for help in making the decision. The doctor is available to meet with all family members together if this might be helpful. Perhaps a discussion with the chaplain or faith leader may help as well.

Tube Feeding Procedure

Placing a PEG tube usually takes about 15 minutes. It involves a number of steps. Liquid food is put into a bag that is delivered into the stomach through this tube.

Tube feeding can be done for a limited amount of time. When the decision is made to place the feeding tube, a decision can also be made that the use of the tube will be reviewed in 1 month or 2 or 3 to see if it is still the right thing to do. If it is felt that the original goals of tube feeding are not met, then a new decision can be made to discontinue the tube feeding.

Alternatives to Tube Feeding

Continuing to feed by mouth (feeding orally) is an option to inserting a PEG. Feeding by mouth also has its advantages and disadvantages. Eating allows a person the ability to enjoy the taste of food and have increased social interaction with others. However, it usually requires a longer period of time to feed someone who has problems eating or swallowing.

Guidelines are intended to be flexible. They serve as reference points or recommendations, not rigid criteria. Guidelines should be followed in most cases, but there is an understanding that, depending on the patient, the setting, the circumstances, or other factors, care can and should be tailored to fit individual needs.

Benefits/Burdens of Tube Feeding/PEG Placement for Adults

Información para pacientes, familias y médicos clínicos Ayuda para tomar decisiones sobre la alimentación por sonda o colocación de sonda de GEP

Dificultad para tragar o comer

Las personas que padecen una enfermedad grave o que tienen una salud frágil a veces tienen dificultad para conseguir los nutrientes que creemos que necesitan para que sus cuerpos funcionen correctamente. Cuando comer o tragar se convierte en una labor difícil, el médico tratará de identificar la causa del problema. Si se pueden realizar tratamientos o ajustes en el entorno para solucionar este problema, el médico se asegurará de tomar las medidas necesarias. Si el problema no se puede solucionar a través de dichas medidas, lo más probable es que el médico hable con el paciente y su familia sobre la posibilidad de usar una sonda de alimentación. Un procedimiento de alimentación por sonda conlleva la colocación de una sonda de GEP (gastrostomía endoscópica percutánea); esto es, una sonda de alimentación que se introduce en el estómago a través de un pequeño orificio en el abdomen.

Conversaciones con el médico sobre la alimentación por sonda

Antes de explicar con detalle la alimentación por sonda, el médico le preguntará al paciente (o a sus seres queridos si el paciente no puede tomar una decisión por sí mismo) si está interesado en el procedimiento de alimentación por sonda. Algunas personas tienen una opinión muy clara sobre la alimentación por sonda y con frecuencia la comparten con sus seres queridos. El médico preguntará si el paciente ha realizado algún tipo de planificación anticipada de cuidados, si tiene un representante para cuidados médicos o si ha redactado un testamento vital. Asimismo, preguntará si el paciente ha hablado alguna vez con sus seres queridos acerca de sus preferencias de asistencia médica en ese tipo de situaciones. Determinar si se desea seguir una alimentación por sonda suele ser una decisión difícil.

Para tomar este tipo de decisión por un ser querido, es importante saber discernir qué es lo que el paciente querría si pudiera tomar dicha decisión por sí mismo, así como ser capaz de separar la voluntad del paciente de lo que uno desea para este. La decisión debería tomarse en función del deseo del paciente.

La decisión sobre la alimentación por sonda

A la hora de tomar una decisión sobre la alimentación por sonda, se han de tener muchos aspectos en cuenta. Es importante considerar las ventajas, desventajas y otras consideraciones en relación con la colocación de la sonda de alimentación. Asimismo, es importante analizar las ventajas, desventajas y otras consideraciones de continuar la alimentación por vía oral.

A la hora de tomar este tipo de decisión, uno puede hacerse las siguientes preguntas:

- ¿La vida de mi ser querido se alargará o se acortará como resultado de la colocación de una sonda de GEP?
- ¿Mejorará su calidad de vida? ¿Valoraría mi ser querido dicha calidad de vida?
- ¿Colocarle una sonda de GEP podría hacer posible la administración de un tratamiento que cure la enfermedad subyacente?

Por ejemplo, la alimentación por sonda dará resultados diferentes en una persona que ha sufrido un derrame cerebral pero que antes disfrutaba de una buena salud que en una persona que tiene la enfermedad de Alzheimer.

Las emociones suelen jugar un papel importante a la hora de decidir si se desea colocar una sonda de alimentación. Los sentimientos de culpa, como “no hacer todo lo que está en el poder de uno” para ayudar al paciente, y la presión de fuentes externas puede afectar al proceso de toma de decisiones. Por último, la opinión personal sobre la alimentación por sonda también influye en la decisión. Los representantes para cuidados médicos y los familiares deben tener en cuenta muchos aspectos antes de tomar una decisión sobre la alimentación por sonda. Las personas que optan por no seguir una alimentación por sonda pueden llevar una vida cómoda ingiriendo líquidos con pequeños sorbos y lubricándose la boca y los labios. La mayoría de los pacientes no sienten una mayor comodidad si se les coloca una sonda de alimentación. Existen algunas excepciones, como pacientes con lesiones graves que merman su capacidad para tragar y personas con cáncer incipiente de cabeza, cuello y esófago.

Si usted y sus familiares no se ponen de acuerdo en si al paciente se le debería colocar una sonda de alimentación o no, es importante pedir ayuda para tomar una decisión. El médico estaría dispuesto a reunirse con todos los miembros de la familia si de este modo pudiera ayudar en la toma de decisiones. Hablar con un capellán o guía religioso también puede resultar de ayuda.

El procedimiento de la alimentación por sonda

La colocación de una sonda de GEP suele llevar alrededor de quince minutos y comprende una serie de pasos. Se colocan alimentos líquidos en una bolsa y se introducen en el estómago a través de la sonda.

La alimentación por sonda puede realizarse durante un periodo de tiempo limitado. Una vez que se ha tomado la decisión de colocar la sonda de alimentación, también se puede decidir que se examine el uso de la sonda tras uno, dos o tres meses de su colocación para ver si sigue siendo la decisión correcta. Si se determina que no se están cumpliendo los objetivos originales de la alimentación por sonda, se puede tomar una nueva decisión para interrumpir el procedimiento.

Alternativas a la alimentación por sonda

Seguir realizando una alimentación por la boca (por vía oral) es una alternativa a la inserción de una sonda de GEP. La alimentación por vía oral también tiene sus ventajas y desventajas. Comer aporta placer al paciente al poder saborear los alimentos y mejora su interacción social con otras personas. Sin embargo, por lo general lleva más tiempo alimentar a personas que tienen dificultades para comer o tragar.

Benefits/Burdens of Tube Feeding/PEG Placement for Adults

**Patient/Family/Clinician Information
Benefits and Burdens of Tube Feeding/PEG Placement**

GOALS FOR CARE

DISEASES	Prolongs Life	Improves Quality of Life or Functional Ability³	Enables a Cure or Reverses the Disease Process
Stroke (good health in general before this)	Likely	Up to 25% regain ability to swallow	Not Likely
Stroke (in poor health before this)	Likely in the short term	Not Likely	Not Likely
	Not likely in the long term		
Neurodegenerative Disease [for example, Amyotrophic Lateral Sclerosis (ALS)] ¹	Likely	Uncertain	Not Likely
Persistent Vegetative State(PVS) ²	Likely	Not Likely	Not Likely
Advanced Organ Failure	Not Likely	Not Likely	Not Likely
Frailty	Not Likely	Not Likely	Not Likely
Advanced Dementia	Not Likely	Not Likely	Not Likely
Advanced Cancer	Not Likely	Not Likely	Not Likely

This information is based predominately on a consensus of current expert opinion. It is not exhaustive. There are always patients who provide exceptions to the rule.

1. A severe disease affecting the brain and spinal cord.
2. Person with severe brain damage with no awareness.
3. Ability to do things like eating, dressing, going to the bathroom without assistance.

Benefits/Burdens of Tube Feeding/PEG Placement for Adults

Hoja informativa para el paciente/familia/el clínico Beneficios y dificultades del Alimentación por tubo/PEG

ETAS PARA EL CUIDADO

ENFERMEDADES	Prolonga la vida	Mejora la calidad de vida o habilidad funcional ³	Permite la cura o revierte el proceso de la enfermedad
Derrame cerebral (buena salud en general antes de esto)	Probable	Posibilidad de recuperar la habilidad de tragar de hasta 25%	Poco probable
Derrame cerebral (salud pobre antes de esto)	Probable a corto plazo	Poco probable	Poco probable
	Poco probable a largo plazo		
Neurodegenerativas Esclerosis Lateral [por ejemplo, esclerosis lateral amiotrófica (EAL-ALS en inglés)] ¹	Probable	Incierto	Poco probable
Estado vegetativo persistente (EVP) ²	Probable	Poco probable	Poco probable
Fallo avanzado de los órganos	Poco probable	Poco probable	Poco probable
Debilidad	Poco probable	Poco probable	Poco probable
Demencia avanzada	Poco probable	Poco probable	Poco probable
Cáncer avanzado	Poco probable	Poco probable	Poco probable

Esta información se basa principalmente en un consenso de la opinión actual de expertos. De ninguna manera es exhaustiva. Siempre hay pacientes cuya experiencia provee excepciones a la regla.

1. Una enfermedad grave que afecta el cerebro o la médula espinal
2. Persona con daño cerebral grave sin conciencia
3. Habilidad de hacer cosas tales como comer, vestirse, ir al baño sin asistencia

Benefits/Burdens of Tube Feeding/PEG Placement for Adults

References

- Anis M, Abid S, Jafri W et al. Acceptability and outcomes of the Percutaneous Endoscopic Gastrostomy (PEG) tube placement – patients’ and care givers’ perspectives. BMC Gastroenterology 2006;6:37.
- Bankhead, R., et al. A.S.P.E.N. Enteral Nutrition Practice Recommendations. Journal of Parenteral Nutrition. 2009;33:163-165. <http://pen.sagepub.com/content/33/2/122.full.pdf>
- Bannerman E, Pendlebury J, Phillips F, Ghosh S. A Cross-Sectional and Longitudinal Study of Health-related Quality of Life after Percutaneous Gastrostomy. Eur J Gastroenterol Hepatol. 2000 Oct;12(10):1101-9.
- Bannerman E, Phillips F, Pendlebury J, Ghosh S. Cross-Sectional and Prospective Studies of Nutritional Indices after Percutaneous Gastrostomy. Eur J Gastroenterol Hepatol. 2001 Nov;13(11):1315-21.
- Barrett JS, Shepherd, SJ, Gibson, PR. Strategies to Manage Gastrointestinal Symptoms Complicating Enteral Feeding. Journal of Parenteral and Enteral Nutrition. 2009;33:21-26. <http://pen.sagepub.com/content/33/1/21.abstract>
- Braun U, Rabeneck, L, McCullough L, et al. Decreasing Use of Percutaneous Endoscopic Gastrostomy Tube Feeding for Veterans with Dementia – Racial Differences Remain. JAGS 2005 53:242-248.
- Brett AS, Dementia, Gastrostomy Tubes, and Mortality. Arch Intern Med. 2001 Oct 22;161(19):2385-6. Brett AS, Rosenberg JC. The Adequacy of Informed Consent for Placement of Gastrostomy Tubes. Arch Intern Med. 2001 Mar 12;161(5):745-8.
- Callahan CM, Haag KM, Weinberger M, Tierney WM, Buchanan NN, Stump TE, Nisi R. Outcomes of Percutaneous Endoscopic Gastrostomy Among Older Adults in a Community Setting. J Am Geriatr Soc. 2000 Sep;48(9):1048-54.
- Callahan CM, Haag KM, Buchanan NN, Nisi R. Decision-making for Percutaneous Endoscopic Gastrostomy Among Older Adults in a Community Setting. J Am Geriatr Soc. 1999 Sept;47(9):1105-9.
- Dennis M. Nutrition After Stroke. Br Med Bull. 2000;56(2):466-75.
- Doig G, Simpson, F, Finfer S. et al. Effect of Evidence-Based Feeding Guidelines on Mortality of Critically Ill Adults: A Cluster Randomized Control Trial. JAMA. 2008;300 (23):2731-2741.
- Dwolatzky T, Berezovski S, Friedmann R, et al. A Prospective Comparison of the Use of Nasogastric and Percutaneous Endoscopic Gastrostomy Tubes for Long-term Enteral Feeding in Older People. Clin Nutr. 2001 Dec 20;(6):535-40.
- Baird EA et al. Early Prediction of Stroke Recovery: A Three-Item Scale for Early Prediction of Stroke Recovery. Lancet 2001;357:2095-2099.
- Elia M, Stratton RJ, Holden C, et al.; Committee of the British Artificial Nutrition Survey (BANS). Home Enteral Tube Feeding Following Cerebrovascular Accident. Clin Nutr. 2001 Feb 20;(1):27-30.
- Finucane TE, Christmas C, Travis K. Tube Feeding in Patients with Advanced Dementia: A Review of the Evidence. JAMA. 1999 Oct 13;282(14):1365-70.
- Finucane TE, Christmas C, Leff BA. Tube feeding in dementia: how incentives undermine health care quality and patient safety. Journal of American Medical Directors Association. 2007;8:205-208. <http://www.ncbi.nlm.nih.gov/pubmed/17498602>
- Genao L, White H, Twersky J. The clinical course of advanced dementia. New England Journal of Medicine. 2010;362:363-364.
- Gessert CE, Mosier MC, Brown EF, et al. Tube feeding in nursing home residents with severe and irreversible cognitive impairment. Journal of American Geriatric Society. 2000;48:1593-1600. <http://www.ncbi.nlm.nih.gov/pubmed/11129748>

Benefits/Burdens of Tube Feeding/PEG Placement for Adults

References, cont'd.

- Grant MD, Rudberg MA, Brody JA. Gastrostomy Placement and Mortality Among Hospitalized Medicare Beneficiaries. JAMA. 1998 Jun 24; 279(24):1973-6.
- Graham I, Logan J, O'Connor A. et al. A Qualitative Study of Physicians Perceptions of Three Decision Aids. Patient Education and Counseling. Patient Educ Couns. 2003 Jul;50(3):279-83 <http://www.ncbi.nlm.nih.gov/pubmed/12900100>
- Hallenbeck J, Weissman D. Fast Fact and Concept #10: Tube Feed or Not Tube Feed? Journal of Palliative Medicine. 2002 Dec;Vol. 5, No. 6: 909-910.
- Holloway R, Benesch C, Burgin WS, Zentner J. Palliative Care in Stroke. Department of Neurology and Department of Community and Preventive Medicine, University of Rochester School of Medicine, Rochester, New York. January 12, 2005 Manuscript.
- James A, Kapur K, Hawthorne. Long-term Outcome of Percutaneous Endoscopic Gastrostomy Feeding in Patients with Dysphagic Stroke. Age and Aging. 1998 Nov;27(6):671-6.
- Johnston D, Tham T, Mason M. Death after PEG: results of the National Confidential Enquiry into Patient Outcome and Death. Gastrointestinal Endoscopy 2008;68 (2):223-227.
- Kobayashi K, Cooper GS, Grant MD, Rudberg MA. Gastrostomy placement and mortality among hospitalized Medicare beneficiaries. Gastrointest Endosc. 1999 Jun;49(6):810-1.
- Lopez RP, Amella EJ, Strumpf NE, et al. The influence of nursing home culture on the use of feeding tubes. Archives of Internal Medicine. 2010;170:83-88.
- Meier DE, Ahronheim JC, Morris J, et al. High short-term mortality in hospitalized patients with advanced dementia: lack of benefit of tube feeding. Archives of Internal Medicine. 2001;161:594-599.
- Mitchell SL, Teno JM, Roy J, Kabumoto G, Mor V. Clinical and Organizational Factors Associated with Feeding Tube Use Among Nursing Home Residents with Advanced Cognitive Impairment. JAMA. 2003 July 2;290(1):73-80.
- Mitchell SL, Buchanan JL, Littlehale S, Hamel MB. Tube-feeding versus hand-feeding nursing home residents with advanced dementia: a cost comparison. J Am Med Dir Assoc. 2003 Jan-Feb;4(1):27-33.
- Mitchell SL. Financial Incentives for Placing Feeding Tubes in Nursing Home Residents with Advanced Dementia. J Am Geriatr Soc. 2003 Jan;51(1):129-31.
- Mitchell SL, Tetroe J, O'Connor AM. A Decision Aid for Long-term Tube Feeding in Cognitively Impaired Older Persons. J Am Geriatr Soc. 2001 Mar;49(3):313-6.
- Mitchell SL, Tetroe JM. Survival After Percutaneous Endoscopic Gastrostomy Placement in Older Persons. J Gerontol A Biol Sci Med Sci. 2000 Dec;55(12): M735-9.
- Mitchell SL, Berkowitz RE, Lawson FM, Lipsitz LA. A Cross-National Survey of Tube-Feeding Decisions in Cognitively Impaired Older Persons. J Am Geriatr Soc. 2000 Apr;48(4):391-7.
- Mitchell SL, Kiely DK, Gillick MR. Nursing home characteristics associated with tube feeding in advanced cognitive impairment. Journal of American Geriatric Society. 2003;51:75-79. <http://www.ncbi.nlm.nih.gov/pubmed/12534849>.
- Modi S, Whetstone L, Cummings D. Influence of Patient and Physician Characteristics on Percutaneous Endoscopic Gastrostomy Tube Decision-Making. Journal of Palliative Medicine. 2007;10 (2):359-366.
- Monteleoni C, Clark E. Using Rapid-Cycle Quality Improvement Methodology to Reduce Feeding Tubes in Patients with Advance Dementia: Before and After Study. BMJ. 2004 Aug 24;(329):491-94.
- Murphy DJ, Santilli S. Elderly patients' preferences for long-term life support. Archives of Family Medicine, 1998;7:484-488.

Benefits/Burdens of Tube Feeding/PEG Placement for Adults

References, cont'd.

- Nair S, Herten H, Pitchumoni CS. Hypoalbuminemia Is a Poor Predictor of Survival after Percutaneous Endoscopic Gastrostomy in Elderly Patients with Dementia. *Am J Gastroenterol.* 2000 Jan;(95)1:133-6.
- Pennington C. To PEG or Not to PEG. *Clin Med.* 2002 May-Jun;2(3):250-5.
- Rabeneck L, McCullough LB, Wray NP. Ethically Justified, Clinically Comprehensive Guidelines for Percutaneous Endoscopic Gastrostomy Tube Placement. *Lancet.* 1997 Feb 15;349(9050):496-8.
- Rabeneck L, Wray NP, Petersen NJ. Long-Term Outcomes of Patients Receiving Percutaneous Endoscopic Gastrostomy Tubes. *J Gen Intern Med.* 1996 May;11(5):287-93.
- Rudberg MA, Egleston BL, Grant MD, Brody JA. Effectiveness of Feeding Tubes in Nursing Home Residents with Swallowing Disorders. *JPEN J Parenter Enteral Nutr.* 2000 Mar-Apr;24(2):97-102.
- Sampson EL, Candy B, Jones L. Enteral tube feeding for older people with advanced dementia. *Cochrane Database of Systematic Reviews* 2009, Issue 2. Art. No.: CD007209. DOI: 10.1002/14651858.CD007209.pub2.
<http://www.ncbi.nlm.nih.gov/pubmed/19370678>.
- Sanders H, Newall S, Norton B, Holmes GT. Gastrostomy Feeding in the Elderly After Acute Dysphagic Stroke. *J Nutr Health Aging.* 2000;4(1):58-60.
- Sanders H, Carter MJ, D'Silva J, James G, Bolton RP, Willemse PJ, Bardhan, KD. Percutaneous Endoscopic Gastrostomy: A Prospective Audit of the Impact of Guidelines in Two District General Hospitals in the United Kingdom. *Am. J Gastroenterol* 2002; 97(9):2239-45.
- Sanders S, Anderson A, et al. Percutaneous Endoscopic Gastrostomy: An Effective Strategy for Gastrostomy Feeding in Patients with Dementia. *Clinical Medicine* May 2004;4(3):235-241.
- Sheiman SL. Tube feeding the demented nursing home resident. *Journal of American Geriatric Society.* 1996;44:1268-1270.
- Skelly RH. Are We Using Percutaneous Endoscopic Gastrostomy Appropriately in the Elderly? *Curr Opin Clin Nutr Metab Care.* 2002 Jan;5(1) 35-42.
- Skelly RH, Kupfer, RM, Metcalfe ME, Allison SP, Holt M, Hull MA, Rawlings JK. Percutaneous Endoscopic Gastrostomy (PEG): Change in Practice Since 1988. *Clin Nutr.* 2002 Oct;21(5):389-94.
- Slomka J. Withholding Nutrition at the End of Life: Clinical and Ethical Issues. *Cleveland Clinic Journal of Medicine* June 2003;70(6): 548-552.
- Smith BM, Perring P, Engoren, M, Sferra J. Hospital and Long-Term Outcome after Percutaneous Endoscopic Gastrostomy. *Surg Endosc.* 2008;22:74-80.
- Taylor PR. Decision Making in Long-Term Care: Feeding Tubes. *Annals of Long-Term Care: Clinical Care Aging.* 2001;9(11): 21-6.
- Teno JM, Gozalo P, Mitchell S, Kuo S, Fulton A, Mor V. Feeding Tubes and the Prevention or Healing of Pressure Ulcers. *Arch Intern Med.* 2012;172(9):697-701.
- Teno JM, Mitchell SL, Gozalo PL, et al. Hospital characteristics associated with feeding tube placement in nursing home residents with advanced cognitive impairment. *Journal of the American Medical Association.* 2010;303:544-550.
- Thomas DR, Ashmen W, Morley JE, Evans, WJ. Nutritional Management in Long-Term Care. *J Gerontology Series A.* 2000;(55) M725-34.
- Wijdicks EF, McMahon MM. Percutaneous Endoscopic Gastrostomy After Acute Stroke: Complications and Outcome. *Cerebrovasc Dis.* 1999 Mar-Apr;9(2):109-11.