Benefits/Burdens of Tube Feeding/PEG Placement for Adults References



Anis M, Abid S, Jafri W et al. Acceptability and outcomes of the Percutaneous Endoscopic Gastrostomy (PEG) tube placement – patients' and care givers' perspectives. BMC Gastroenterology 2006; 6:37.

Bankhead, R., et al. A.S.P.E.N. Enteral Nutrition Practice Recommendations. Journal of Parenteral Nutrition. 2009; 33:163-165. http://pen.sagepub.com/content/33/2/122.full.pdf

Bannerman E, Pendlebury J, Phillips F, Ghosh S. A Cross-Sectional and Longitudinal Study of Health-related Quality of Life after Percutaneous Gastrostomy. Eur J Gastroenterol Hepatol. 2000 Oct;12(10):1101-9.

Bannerman E, Phillips F, Pendlebury J, Ghosh S. Cross-Sectional and Prospective Studies of Nutritional Indices after Percutaneous Gastrostomy. Eur J Gastroenterol Hepatol. 2001 Nov;13(11):1315-21.

Barrett JS, Shepherd, SJ, Gibson, PR. Strategies to Manage Gastrointestinal Symptoms Complicating Enteral Feeding. Journal of Parenteral and Enteral Nutrition. 2009; 33:21-26. http://pen.sagepub.com/content/33/1/21.abstract

Braun U. Rabeneck, L, McCullough L, et al. Decreasing Use of Percutaneous Endoscopic Gastrostomy Tube Feeding for Veterans with Dementia – Racial Differences Remain. JAGS 2005 53:242-248.

Brett AS, Dementia, Gastrostomy Tubes, and Mortality. Arch Intern Med. 2001 Oct 22;161(19):2385-6. Brett AS, Rosenberg JC. The Adequacy of Informed Consent for Placement of Gastrostomy Tubes. Arch Intern Med. 2001 Mar 12;161(5):745-8.

Callahan CM, Haag KM, Weinberger M, Tierney WM, Buchanan NN, Stump TE, Nisi R. Outcomes of Percutaneous Endoscopic Gastrostomy Among Older Adults in a Community Setting. J Am Geriatr Soc. 2000 Sep;48(9):1048-54.

Callahan CM, Haag KM, Buchanan NN, Nisi R. Decision-making for Percutaneous Endoscopic Gastrostomy Among Older Adults in a Community Setting. J Am Geriatr Soc. 1999 Sept;47(9):1105-9.

Dennis M. Nutrition After Stroke. Br Med Bull. 2000;56(2):466-75.

Doig G, Simpson, F, Finfer S. et al. Effect of Evidence-Based Feeding Guidelines on Mortality of Critically III Adults: A Cluster Randomized Control Trial. JAMA. 2008;300 (23):2731-2741.

Dwolatzky T, Berezovski S, Friedmann R, et al. A Prospective Comparison of the Use of Nasogastric and Percutaneous Endoscopic Gastrostomy Tubes for Long-term Enteral Feeding in Older People. Clin Nutr. 2001 Dec 20;(6):535-40.

Baird EA et al. Early Prediction of Stroke Recovery: A Three-Item Scale for Early Prediction of Stroke Recovery. Lancet 2001; 357:2095-2099.

Elia M, Stratton RJ, Holden C, et al.; Committee of the British Artificial Nutrition Survey (BANS). Home Enteral Tube Feeding Following Cerebrovascular Accident. Clin Nutr. 2001 Feb 20;(1):27-30.

Finucane TE, Christmas C, Travis K. Tube Feeding in Patients with Advanced Dementia: A Review of the Evidence. JAMA. 1999 Oct 13;282(14):1365-70.

Finucane TE, Christmas C, Leff BA. Tube feeding in dementia: how incentives undermine health care quality and patient safety. Journal of American Medical Directors Association. 2007; 8:205-208. http://www.ncbi.nlm.nih.gov/pubmed/17498602

Genao L, White H, Twersky J. The clinical course of advanced dementia. New England Journal of Medicine. 2010; 362:363-364.

Gessert CE, Mosier MC, Brown EF, et al. Tube feeding in nursing home residents with severe and irreversible cognitive impairment. Journal of American Geriatric Society. 2000; 48:1593-1600. http://www.ncbi.nlm.nih.gov/pubmed/11129748

- Guidelines are intended to be flexible. They serve as reference points or recommendations, not rigid criteria. Guidelines should be followed in most cases, but there is an understanding that, depending on the patient, the setting, the circumstances, or other factors, care can and should be tailored to fit individual needs. For patients associated with OPWDD a separate process must be followed, see OPWDD checklists.

Benefits/Burdens of Tube Feeding/PEG Placement for Adults References, cont'd.



Grant MD, Rudberg MA, Brody JA. Gastrostomy Placement and Mortality Among Hospitalized Medicare Beneficiaries. JAMA. 1998 Jun 24; 279(24):1973-6.

Graham I, Logan J, O'Connor A. et al. A Qualitative Study of Physicians Perceptions of Three Decision Aids. Patient Education and Counseling. Patient Educ Couns. 2003 Jul;50(3):279-83 http://www.ncbi.nlm.nih.gov/pubmed/12900100

Hallenbeck J, Weissman D. Fast Fact and Concept #10: Tube Feed or Not Tube Feed? Journal of Palliative Medicine. 2002 Dec; Vol. 5, No. 6: 909-910.

Holloway R, Benesch C, Burgin WS, Zentner J. Palliative Care in Stroke. Department of Neurology and Department of Community and Preventive Medicine, University of Rochester School of Medicine, Rochester, New York. January 12, 2005 Manuscript.

James A, Kapur K, Hawthorne. Long-term Outcome of Percutaneous Endoscopic Gastrostomy Feeding in Patients with Dysphagic Stroke. Age and Aging. 1998 Nov;27(6):671-6.

Johnston D, Tham T, Mason M. Death after PEG: results of the National Confidential Enquiry into Patient Outcome and Death. Gastrointestinal Endoscopy 2008;68 (2):223-227.

Kobayashi K, Cooper GS, Grant MD, Rudberg MA. Gastrostomy placement and mortality among hospitalized Medicare beneficiaries. Gastrointest Endosc. 1999 Jun;49(6):810-1.

Lopez RP, Amella EJ, Strumpf NE, et al. The influence of nursing home culture on the use of feeding tubes. Archives of Internal Medicine. 2010; 170:83-88.

Meier DE, Ahronheim JC, Morris J, et al. High short-term mortality in hospitalized patients with advanced dementia: lack of benefit of tube feeding. Archives of Internal Medicine. 2001; 161:594-599.

Mitchell SL, Teno JM, Roy J, Kabumoto G, Mor V. Clinical and Organizational Factors Associated with Feeding Tube Use Among Nursing Home Residents with Advanced Cognitive Impairment. JAMA. 2003 July 2;290(1):73-80.

Mitchell SL, Buchanan JL, Littlehale S, Hamel MB. Tube-feeding versus hand-feeding nursing home residents with advanced dementia: a cost comparison. J Am Med Dir Assoc. 2003 Jan-Feb;4(1):27-33.

Mitchell SL. Financial Incentives for Placing Feeding Tubes in Nursing Home Residents with Advanced Dementia. J Am Geriatr Soc. 2003 Jan;51(1):129-31.

Mitchell SL, Tetroe J, O'Connor AM. A Decision Aid for Long-term Tube Feeding in Cognitively Impaired Older Persons. J Am Geriatr Soc. 2001 Mar;49(3):313-6.

Mitchell SL, Tetroe JM. Survival After Percutaneous Endoscopic Gastrostomy Placement in Older Persons. J Gerontol A Biol Sci Med Sci. 2000 Dec;55(12): M735-9.

Mitchell SL, Berkowitz RE, Lawson FM, Lipsitz LA. A Cross-National Survey of Tube-Feeding Decisions in Cognitively Impaired Older Persons. J Am Geriatr Soc. 2000 Apr;48(4):391-7.

Mitchell SL, Kiely DK, Gillick MR. Nursing home characteristics associated with tube feeding in advanced cognitive impairment. Journal of American Geriatric Society. 2003; 51:75-79. http://www.ncbi.nlm.nih.gov/pubmed/12534849.

Modi S, Whetstone L, Cummings D. Influence of Patient and Physician Characteristics on Percutaneous Endoscopic Gastrostomy Tube Decision-Making. Journal of Palliative Medicine. 2007;10 (2):359-366.

Monteleoni C, Clark E. Using Rapid-Cycle Quality Improvement Methodology to Reduce Feeding Tubes in Patients with Advance Dementia: Before and After Study. BMJ. 2004 Aug 24;(329):491-94.

Murphy DJ, Santilli S. Elderly patients' preferences for long-term life support. Archives of Family Medicine, 1998;7:484-488.

- Guidelines are intended to be flexible. They serve as reference points or recommendations, not rigid criteria. Guidelines should be followed in most cases, but there is an understanding that, depending on the patient, the setting, the circumstances, or other factors, care can and should be tailored to fit individual needs. For patients associated with OPWDD a separate process must be followed, see OPWDD checklists.

Benefits/Burdens of Tube Feeding/PEG Placement for Adults



References, cont'd.

Nair S, Hertan H, Pitchumoni CS. Hypoalbuminemia Is a Poor Predictor of Survival after Percutaneous Endoscopic Gastrostomy in Elderly Patients with Dementia. Am J Gastroenterol. 2000 Jan; (95) 1:133-6.

Pennington C. To PEG or Not to PEG. Clin Med. 2002 May-Jun;2(3):250-5.

Rabeneck L, McCullough LB, Wray NP. Ethically Justified, Clinically Comprehensive Guidelines for Percutaneous Endoscopic Gastrostomy Tube Placement. Lancet. 1997 Feb 15;349(9050):496-8.

Rabeneck L, Wray NP, Petersen NJ. Long-Term Outcomes of Patients Receiving Percutaneous Endoscopic Gastrostomy Tubes. J Gen Intern Med. 1996 May;11(5):287-93.

Rudberg MA, Egleston BL, Grant MD, Brody JA. Effectiveness of Feeding Tubes in Nursing Home Residents with Swallowing Disorders. JPEN J Parenter Enteral Nutr. 2000 Mar-Apr;24(2):97-102.

Sachdev, S., Refaat, T., Bacchus, I.D. et al. Age most significant predictor of requiring enteral feeding in head-and-neck cancer patients. Radiat Oncol 10, 93 (2015).

Sachdev, S., Refaat, T., Bacchus, I.D. et al. Age most significant predictor of requiring enteral feeding in head-and-neck cancer patients. Radiat Oncol 10, 93 (2015). https://rdcu.be/b2MvX

Sampson EL, Candy B, Jones L. Enteral tube feeding for older people with advanced dementia. Cochrane Database of Systematic Reviews 2009, Issue 2. Art. No.: CD007209. DOI: 10.1002/14651858.CD007209.pub2. http://www.ncbi.nlm.nih.gov/pubmed/19370678.

Sanders H, Newall S, Norton B, Holmes GT. Gastrostomy Feeding in the Elderly After Acute Dysphagic Stroke. J Nutr Health Aging. 2000;4(1):58-60.

Sanders H, Carter MJ, D'Silva J, James G, Bolton RP, Willemse PJ, Bardhan, KD. Percutaneous Endoscopic Gastrostomy: A Prospective Audit of the Impact of Guidelines in Two District General Hospitals in the United Kingdom. Am. J Gastroenterol 2002; 97(9):2239-45.

Sanders S, Anderson A, et al. Percutaneous Endoscopic Gastrostomy: An Effective Strategy for Gastrostomy Feeding in Patients with Dementia. Clinical Medicine May 2004;4(3):235-241.

Sheiman SL. Tube feeding the demented nursing home resident. Journal of American Geriatric Society. 1996; 44:1268-

1270. Skelly RH. Are We Using Percutaneous Endoscopic Gastrostomy Appropriately in the Elderly? Curr Opin Clin Nutr

Metab

Care. 2002 Jan;5(1) 35-42.

Skelly RH, Kupfer, RM, Metcalfe ME, Allison SP, Holt M, Hull MA, Rawlings JK. Percutaneous Endoscopic Gastrostomy (PEG): Change in Practice Since 1988. Clin Nutr. 2002 Oct;21(5):389-94.

Slomka J. Withholding Nutrition at the End of Life: Clinical and Ethical Issues. Cleveland Clinic Journal of Medicine June 2003;70(6): 548-552.

Smith BM, Perring P, Engoren, M, Sferra J. Hospital and Long-Term Outcome after Percutaneous Endoscopic Gastrostomy. Surg Endosc. 2008; 22:74-80.

Taylor PR. Decision Making in Long-Term Care: Feeding Tubes. Annals of Long-Term Care: Clinical Care Aging. 2001;9(11): 21-6.

Teno JM, Gozalo P, Mitchell S, Kuo S, Fulton A, Mor V. Feeding Tubes and the Prevention or Healing of Pressure Ulcers. Arch Intern Med. 2012;172(9):697-701.

Teno JM, Mitchell SL, Gozalo PL, et al. Hospital characteristics associated with feeding tube placement in nursing home residents with advanced cognitive impairment. Journal of the American Medical Association. 2010; 303:544-550.

Thomas DR, Ashmen W, Morley JE, Evans, WJ. Nutritional Management in Long-Term Care. J Gerontology Series A. 2000;(55) M725-34.